GENERAL VASCULAR SURGERY MEDICAL GROUP, INC.

Phone: (510) 357-4006 Fax: (510) 347-4712

ROBERT O. GINGERY, M.D. ARNOLD W. LEVINE, M.D. LAMONT D. PAXTON, M.D. MICHAEL D. INGEGNO, M.D. DONATO J. STINGHEN VASCULAR SURGERY NONINVASIVE VASCULAR DIAGNOSIS

POST-OPERATIVE INFORMATION following CAROTID ENDARTERECTOMY

1	3.T 1		•	C 1:
1	Normal	post-operat	111/10	tindings.
1.	Month	post operat	11 4 C	mumas.

Mild swelling; may feel like a hard knot along incision

Area of numbness around your incision; may involve the ear, neck, and part of the face. If you have incisional pain, you may use Tylenol. You may take up to 2 tablets every four hours as needed for pain.

- 2. If at any time following your surgery you have symptoms of a stroke or slight stroke, such as an episode of blindness in on eye, slurred speech, numbness, or weakness in an arm or leg, you should notify your doctor immediately.
- 3. Following discharge from the hospital, you should call your doctor's office and make an appointment to be seen in the office within 7-10 days.
- 4. Your doctor does not want you to drive a motor vehicle until after you have been seen in the office and he gives you permission.
- 5. Your incision should be kept dry and clean until you are seen for your first post-operative office visit.
- 6. Dissolvable sutures were used to close your incision. These will dissolve on their own and do not have to be removed by your doctor.
- 7. You are to continue taking the same medication at the same dosages you were taking prior to surgery unless instructed otherwise by your doctor.
- 8. Walking is good for you. We encourage it daily. Just remember you will tire easily, so take it slowly at first and gradually increase your activity.
- 9. If you have any questions, please feel free to call our office at the above-listed number.